

Matrix Theory and Feynman Diagrams¹

J. Plefka¹, M. Serone² and A. Waldron³

¹ Albert-Einstein-Institut

Max-Planck-Institut für Gravitationsphysik

Schlaatzweg 1, 14473 Potsdam, Germany

² Department of Mathematics, University of Amsterdam

Plantage Muidergracht 24, 1018 TV Amsterdam

The Netherlands

³ NIKHEF, P.O. Box 41882, 1009 DB Amsterdam

The Netherlands

Abstract: We briefly review the computation of graviton and antisymmetric tensor scattering amplitudes in Matrix Theory from a diagrammatic S-Matrix point of view.

It is by now commonly believed that eleven dimensional supergravity is the low-energy effective theory of a more fundamental microscopic model, known as M-theory. A proposed definition of M-theory has been given in terms of the large N limit of a quantum mechanical supersymmetric $U(N)$ Yang-Mills model called Matrix Theory [1]. Since the time of this conjecture many computations have been performed to test the proposal [2]. We here wish to focus on the particle like excitations of Matrix Theory, namely the 11-dim supergravity multiplet, and review a setup which enables one to compute various S-Matrix elements amongst them [3, 4]. These amplitudes are then shown to agree with the corresponding tree-level amplitudes of the low energy supergravity theory, thus demonstrating that Matrix Theory indeed has D=11 supergravity as its low energy limit. We emphasize that it is possible to perform this comparison on the familiar level of Feynman diagram amplitudes in momentum space, in particular there is no need to revert to the calculation of effective source-probe potentials on the supergravity side once one employs our definition of the Matrix Theory S-Matrix. This approach appears to be mandatory once one intends to go beyond the tree-level (classical) approximation to study quantum corrections to the effective M-theory action. After all this is the true challenge for Matrix Theory, as the four and six point tree-level amplitudes are believed to be determined by the large amount of supersymmetry in the problem [5].

1 The Matrix Theory S-Matrix

The M-theory matrix model emerges as the reduction of 9+1 dimensional supersymmetric $U(N)$ Yang-Mills to 0+1 dimensions. In euclidean time the quantum mechanical action

¹Talk given by J. Plefka at the 32nd Symposium Ahrenshoop on the Theory of Elementary Particles Buckow, September 1 - 5, 1998.

reads

$$S = \int dt \left[\frac{1}{2} \text{tr}(D_t X^a)^2 - \frac{1}{4} \text{tr}[X^a, X^b]^2 + \frac{1}{2} \text{tr}(\psi^T D_t \psi - \psi^T \gamma_a [X^a, \psi]) \right], \quad (1.1)$$

where $D_t X^a = \partial_t X^a - i[A, X^a]$ and $D_t \psi = \partial_t \psi - i[A, \psi]$, A , X^a and ψ_α are hermitian $U(N)$ matrices, ($a = 1, \dots, 9$ and $\alpha = 1, \dots, 16$). Moreover we employ a real symmetric representation for the Dirac matrices γ_a in which the charge conjugation matrix $\mathcal{C}$ equals unity. One proceeds to compute the background field effective action by expanding the matrices X_{ij}^a and ψ_{ij} ($i = 1, \dots, N$) around the diagonal background

$$X_{ij}^a = \delta_{ij}(v_i^a t + b_i^a) + Y_{ij}^a \quad \psi_{ij} = \delta_{ij}\theta_i + \tilde{\psi}_{ij} \quad (1.2)$$

with the constant velocities v_i , impact paramters b_i and spins θ_i . This background manifestly solves the classical equations of motion. After fixing a standard background field gauge the quantum fields Y^a , $\tilde{\psi}$ and A , as well as the ghost matrices c and $\bar{c}$ are then integrated out of the path integral via a perturbative loop expansion. One finally arrives at the effective background field action taking the form

$$\int_{-T/2}^{T/2} dt \Gamma = \frac{T}{2} \sum_i v_i^2 + \int_{-T/2}^{T/2} dt \sum_{l \geq 1} \Gamma^{(l)}(v_{ij}, r_{ij}(t), \theta_{ij}) \quad (1.3)$$

Note that in the l loop effective potential $\Gamma^{(l)}$ only the relative quantities $v_{ij} = v_i - v_j$, $r_{ij}(t) = v_{ij}t + b_{ij}$ and $\theta_{ij} = \theta_i - \theta_j$ enter, T is the interaction time.

Let us from now on concentrate on the $U(2)$ model being relevant to two body interactions. At one loop the effective potential reads $\Gamma^{(1)} = v_{12}^4/r_{12}(t)^7 + (\theta_{12} \text{-terms})$. In order to compute the corresponding 4-point S-Matrix of Matrix Theory, we prepare two incoming and two outgoing states carrying the quantum numbers momentum p and polarization $\mathcal{H}$ of the four particles involved in the scattering process. The S-Matrix element is then written as

$$\mathcal{S}_{2 \rightarrow 2} = \langle \mathcal{H}^{1'}, \mathcal{H}^{2'} | \int \left(\prod_{i=1}^2 d^9(x'_i, x_i) e^{-ip_i \cdot x_i + ip'_i \cdot x'_i} \exp(i \frac{T}{2} (v_1^2 + v_2^2) + i \int_{-T/2}^{T/2} dt \Gamma^{(1)}) \right) | \mathcal{H}^1, \mathcal{H}^2 \rangle \quad (1.4)$$

We will return shortly to the precise nature of the polarization states $|\mathcal{H}\rangle$. In the above integral one now changes variables to $v_i = (x'_i - x_i)/T$ and $b_i = (x'_i + x_i)/2$ ($i = 1, 2$), and performs the emerging v_i integrals via stationary phase in the relevant $T \rightarrow \infty$ limit. This yields the relation

$$v_i^a = \frac{p_i^{a'} + p_i^a}{2}. \quad (1.5)$$

Expanding out the exponential $\exp(\int dt \Gamma^{(1)})$ and performing the t and b_i integrals finally leads to the matrix element

$$\begin{aligned} \mathcal{S}_{2 \rightarrow 2} = & \delta^9(q_1^a + q_2^a) \delta(q_1 \cdot v_1 + q_2 \cdot v_2) \times_{\theta_1} \langle \mathcal{H}^{1'} |_{\theta_2} \langle \mathcal{H}^{2'} | \left[v_{12}^4 + 2 v_{12}^2 (\theta_{12} \not{v}_{12} \not{v}_{12} \theta_{12}) \right. \\ & \left. + 2 (\theta_{12} \not{v}_{12} \not{v}_{12} \theta_{12})^2 + \frac{4}{9} (\theta_{12} \not{v}_{12} \not{v}_{12} \theta_{12}) (\theta_{12} \not{v}_{12} \gamma^a \theta_{12})^2 + \frac{2}{63} ((\theta_{12} \not{v}_{12} \gamma^a \theta_{12})^2)^2 \right] \frac{1}{q_1^2} | \mathcal{H}^1 \rangle_{\theta_1} \mathcal{H}^2 \rangle_{\theta_2} \end{aligned} \quad (1.6)$$

with the momentum transfer $q_i^a = p_i^{a'} - p_i^a$. Here we have inserted the complete form of the $U(2)$ effective potential, including all spin dependent terms [6, 7]. Note the emergence of the momentum and energy conserving δ -functions in (1.6). In (1.6) the $\theta_{12} = \theta_1 - \theta_2$ are to be interpreted as operators, obeying the algebra $\{\theta_i^\alpha, \theta_j^\beta\} = \delta^{\alpha\beta} \delta_{ij}$. The polarization states

form a representation of this algebra representing the 256 states of the d=11 supergraviton multiplet, i.e.

$$|\mathcal{H}^1\rangle = \mathcal{H}_{\mathcal{M}}^1 |-\rangle_{\theta_1}^{\mathcal{M}}, \quad |\mathcal{H}^2\rangle = \mathcal{H}_{\mathcal{M}}^2 |-\rangle_{\theta_2}^{\mathcal{M}} \quad (1.7)$$

where the generalized index $\mathcal{M}$ denotes $\mathcal{M} \equiv \{ab; abc; ab\alpha\}$, corresponding to the graviton, three-form and gravitino, respectively [8]. These states are built from the diagonal entries θ_1 and θ_2 , contrary to the effective potential in (1.6) depending only on $\theta_1 - \theta_2$. However, via a Fierz rearrangement we may always convert to fermion bilinears homogeneous in θ_1 or θ_2 , as long as we restrict our attention to amplitudes involving bosons only. The general case is a little bit more involved. Whereas $\theta_1 \gamma^{ab} \theta_1$ just acts as a $SO(9)$ rotation on $|-\rangle_{\theta_1}^{ab}$ and $|-\rangle_{\theta_1}^{abc}$, the three index operator acts as

$$\theta_1 \gamma^{mnp} \theta_1 |-\rangle_{\theta_1}^{tu} = -i24\sqrt{3}(\delta^{mt} |-\rangle_{\theta_1}^{npu} - \frac{1}{9} \delta^{tu} |-\rangle_{\theta_1}^{mnp}), \quad (1.8)$$

$$\theta_1 \gamma^{mnp} \theta_1 |-\rangle_{\theta_1}^{tuv} = i24\sqrt{3} \delta^{mt} \delta^{nu} |-\rangle_{\theta_1}^{pv} + \frac{2i}{3} \epsilon^{mnp tuvwyz} |-\rangle_{\theta_1}^{wyz}. \quad (1.9)$$


On the right hand side of (1.8) and (1.9) one must (anti)symmetrize (with unit weight) over all indices according to the symmetry properties of the left hand sides of these equations. Given the algebra of (1.8) and (1.9) only a moderate amount of computer algebra is now required to obtain any pure bosonic S-matrix element. We have computed the 4-graviton amplitude [3] consisting of 66 terms as well as the 4-three form amplitude [4] made out of 103 terms. Due to lack of space we do not state the resulting amplitudes here.

2 The Supergravity Computation


The bosonic sector of 11-dimensional supergravity [9] is given by

$$\begin{aligned} \mathcal{L} = & -\frac{1}{2\kappa^2} \sqrt{-g} R - \frac{1}{8} \sqrt{-g} (F_{MNPQ})^2 \\ & - \frac{\sqrt{3}}{12^3 \kappa} \varepsilon^{M_1 \dots M_{11}} F_{M_1 M_2 M_3 M_4} F_{M_5 M_6 M_7 M_8} C_{M_9 M_{10} M_{11}} \end{aligned} \quad (2.10)$$

where $F_{MNPQ} = 4\partial_{[M} C_{NPQ]}$, $g = \det g_{MN}$ and $M = 0, \dots, 10$. Perturbative quantum gravity is studied by considering small fluctuations h_{MN} from the flat metric η_{MN} , i.e. $g_{MN} = \eta_{MN} + \kappa h_{MN}$ where κ is the 11-dimensional gravitational coupling constant. After employing (for example) the harmonic gauge $\partial_N h^N_M - (1/2)\partial_M h^N_N = 0$ for the graviton as well as the gauge $\partial_M C^M_{NP} = 0$ for the antisymmetric tensor one derives the bosonic propagators in a straightforward fashion. The relevant Feynman diagrams for the two amplitudes considered are the $t = -2p^1 \cdot p^{1'}$ channel ones, which dominate eikonal physics. They are:



$4-g_{MN}$



$4-B_{MNP}$



$4-B_{MNP}$

In order to make a comparison with the Matrix Theory results we need to rewrite our supergravity t channel amplitudes in terms of physical transverse nine dimensional degrees of freedom. Since we are considering the $N = 2$ discrete light cone quantisation formulation [10] of the theory, we work in light cone coordinates and specialise to the case

of vanishing p^- momentum exchange. The kinematics of our $1 + 2 \longrightarrow 1' + 2'$ process then reads

$$\begin{aligned} p_1^M &= (-\tfrac{1}{2} (v_1^a - q_1^a/2)^2, 1, v_1^a - q_1^a/2) & p_{1'}^M &= (-\tfrac{1}{2} (v_1^a + q_1^a/2)^2, 1, v_1^a + q_1^a/2) \\ p_2^M &= (-\tfrac{1}{2} (v_2^a - q_2^a/2)^2, 1, v_2^a - q_2^a/2) & p_{2'}^M &= (-\tfrac{1}{2} (v_2^a + q_2^a/2)^2, 1, v_2^a + q_2^a/2) \end{aligned} \quad (2.11)$$

in units of the compactified radius R and where the 11-dim index M decomposes as $M = (+, -, a)$. We then reduce to physical polarizations by setting $h_{+M} = 0 = h_a^+$, $h_{-M} = -p^a h_{aM}$ as well as $C_{+-a} = 0 = C_{+ab}$ and $C_{-ab} = -\frac{1}{p^-} p_c C_{cab}$. As an example the resulting four graviton amplitude up to terms of order v_{12}^2 reads

$$\begin{aligned} \mathcal{A} &= \frac{1}{q_1^2} \left\{ \frac{1}{2} (h_1 h'_1) (h_2 h'_2) v_{12}^4 + 2 \left[(q_1 h'_2 h_2 v_{12}) (h_1 h'_1) - (q_1 h_2 h'_2 v_{12}) (h_1 h'_1) \right] v_{12}^2 \right. \\ &\quad + (v_{12} h_2 v_{12}) (q_1 h'_2 q_1) (h_1 h'_1) + (v_{12} h'_2 v_{12}) (q_1 h_2 q_1) (h_1 h'_1) - 2 (q_1 h_2 v_{12}) (q_1 h'_2 v_{12}) (h_1 h'_1) \\ &\quad - 2 (q_1 h_1 h'_1 v_{12}) (q_1 h'_2 h_2 v_{12}) + (q_1 h_1 h'_1 v_{12}) (q_1 h_2 h'_2 v_{12}) + (q_1 h'_1 h_1 v_{12}) (q_1 h'_2 h_2 v_{12}) \\ &\quad + \tfrac{1}{2} \left[(q_1 h_1 h'_1 h'_2 h_2 q_1) - 2 (q_1 h_1 h'_1 h_2 h'_2 q_1) + (q_1 h'_1 h_1 h_2 h'_2 q_1) - 2 (q_1 h_2 h'_2 q_1) (h_1 h'_1) \right] v_{12}^2 \\ &\quad \left. + \dots \right\} + [h_1 \longleftrightarrow h_2, h'_2 \longleftrightarrow h'_1] \end{aligned} \quad (2.12)$$

and agrees with the Matrix Theory result. The same holds true for the four three-form amplitude.

References

- [1] T. Banks, W. Fischler, S. Shenker and L. Susskind, Phys. Rev. **D55** (1997) 112, hep-th/9610043.
- [2] K. Becker, M. Becker, J. Polchinski and A. Tseytlin, Phys. Rev. **D65** (1997) 85, hep-th/9706072;
D. Kabat and W. Taylor, Phys. Lett. **B426** (1998) 297, hep-th/9712185;
Y. Okawa and T. Yoneya, Nucl. Phys. **B538** (1999) 67, hep-th/9806108; Nucl. Phys. **B541** (1999) 163, hep-th/9808188.
- [3] J. Plefka, M. Serone and A. Waldron, Phys. Rev. Lett. **81** (1998) 2866.
- [4] J. Plefka, M. Serone and A. Waldron, JHEP **11** (1998) 010.
- [5] S. Paban, S. Sethi and M. Stern, Nucl. Phys. **B534** (1998) 137, hep-th/9805018; JHEP **06** (1998) 012, hep-th/9806028; S. Sethi and M. Stern, hep-th/9903049.
- [6] J.F. Morales, C.A. Scrucca and M. Serone, Phys. Lett. **B417** (1998) 233, hep-th/9709063; Nucl. Phys. **B534** (1998) 223, hep-th/9801183.
- [7] P. Kraus, Phys. Lett. **B419** (1998) 73, hep-th/9709199;
I.N. McArthur, Nucl. Phys. **B534** (1998) 183, hep-th/9806082;
M. Barrio, R. Helling and G. Polhemus, JHEP **05** (1998) 012, hep-th/9801189.
- [8] J. Plefka and A. Waldron, Nucl. Phys. **B512** (1998) 460, hep-th/9710104.
- [9] E. Cremmer, B. Julia and J. Scherk, Phys. Lett. **76B** (1978) 409.
- [10] L. Susskind, *Another conjecture about M(atrix) theory*, hep-th/9704080.